


The Silent Treatment? Changes in patient emotional expression after silence

Christina S. Soma¹  | Bruce E. Wampold² | Nikolaos Flemotomos³ | Raghuveer Peri³ | Shrikanth Narayanan³ | David C. Atkins⁴ | Zac E. Imel¹

¹University of Utah, Salt Lake City, Utah, USA

²University of Wisconsin – Madison, Madison, Wisconsin, USA

³University of Southern California, Los Angeles, California, USA

⁴University of Washington, Seattle, Washington, USA

Correspondence

Christina S. Soma, University of Utah, Salt Lake City, Utah, USA.

Email: tsoma15@gmail.com

Funding information

National Institute on Alcohol Abuse and Alcoholism, Grant/Award Number: R01/AA018673; National Institute on Drug Abuse, Grant/Award Number: K02/AA023814 and R34/DA034860

Abstract

Psychotherapy can be an emotionally laden conversation, where both verbal and non-verbal interventions may impact the therapeutic process. Prior research has postulated mixed results regarding how clients emotionally react following a silence after the therapist is finished talking, potentially due to studying a limited range of silences with primarily qualitative and self-report methodologies. A quantitative exploration may illuminate new findings. Utilising research and automatic data processing from the field of linguistics, we analysed the full range of silence lengths (0.2–24.01 s), and measures of emotional expression—vocally encoded arousal and emotional valence from the words spoken—of 84 audio recordings of Motivational Interviewing sessions. We hypothesised that both the level and the variance of client emotional expression would change as a function of silence length; however, due to the mixed results in the literature, the direction of emotional change was unclear. We conducted a multilevel linear regression to examine how the level of client emotional expression changed across silence length, and an ANOVA to examine the variability of client emotional expression across silence lengths. Results of both analyses indicated that as silence length increased, emotional expression largely remained the same. Broadly, we demonstrated a weak connection between silence length and emotional expression, indicating no persuasive evidence that silence leads to client emotional processing and expression.

KEYWORDS

emotional process, psychotherapy research, linguistics, vocal features, silence

1 | INTRODUCTION

'Silence can exist without speech, but speech cannot exist without silence'. Max Picard (1952).

Psychotherapy is essentially a health-promoting conversation between a therapist and a client (Frank, 1961). Though the words themselves represent the primary vehicle for therapeutic interventions

(Lambert & Bergen, 2004), silence may play an integral part in the therapeutic process (Levitt, 2001a; Levitt, 2002; Frankel et al., 2006; Hill et al., 2003; Daniel et al., 2018; Hill et al., 2019). Silence is not merely a by-product of the conversation (Johannesen, 1974; Levitt, 2002), but a potentially important nonverbal intervention (e.g. Gladstein, 1974).

Psychotherapy researchers have investigated the perceived function and outcome of silences during therapy. Qualitative studies

indicate therapists report using silence to personally reflect on therapeutic content (Hill et al., 2003), as well as a therapeutic intervention (e.g. psychoanalysis, see Freud, 1912; psychodynamic, for a review, see Lane et al., 2002; Hill et al., 2003; Gale & Sanchez, 2005). Silence may be used by the therapist to promote components of the therapeutic relationship, such as rapport building (Sharpley et al., 2005; Daniel et al., 2018), collaboration (Hill et al., 2019; Cuttler et al., 2019) and empathy (Matarazzo & Wiens, 1977; Hill et al., 2003; Ladany et al., 2004). Clients may similarly utilise silence intentionally to formulate their next thought and to reflect on therapeutic content (Levitt, 2001a; Hill et al., 2003). Clients have reported improved attachment to their therapist and decreased feelings of distress after the therapist's use of silence as a therapeutic intervention (Daniel et al., 2018). However, client silences have also been perceived as resistance (e.g. Freud, 1912; Lane et al., 2002). Frequent silences from clients may also be an attempt to disengage with the therapy process, potentially compounding existing insecure attachment and poorer alliance with their therapists (Daniel et al., 2018). As a therapeutic process component, silence may have many uses and consequences depending on contextual factors.

Therapists may specifically use silence in an attempt to facilitate therapeutic tasks, such as processing emotions (Ladany et al., 2004; see Greenberg & Safran, 1989, for a review on emotions in therapy). Silence might correspond with changes in client emotional expression throughout a session (Levitt, 2001b). Qualitative research has indicated therapists use silence to create a space for their clients to experience a particular emotion (Hill et al., 2003). Clients, in turn, have reported using silence to sit and process their emotions (Levitt, 2001b). Silences the client perceives to be helpful could contribute to decreased feelings of overall distress (Frankel et al., 2006) and facilitate an understanding of self or insight into personal experiences (Hill et al., 2019; Levitt, 2001a). Emotion processing is a critical component of psychotherapy process (Greenberg & Safran, 1989) and outcome, including, but not limited to, forming the therapeutic alliance (e.g. Safran & Muran, 2000), client decision-making (Bar-On et al., 2004; Isen, 2008) and physiological changes such as pupil dilation (Lang & Bradley, 2010), heart rate increases (e.g. Helm et al., 2014) and changes in the immune system (e.g. Ackerman et al., 1998; Kiecolt-Glaser et al., 2005).

In addition to silence supporting clients engaging emotionally in therapy, clients have also reported using silence to deliberately disengage from their emotions (Frankel & Levitt, 2009), potentially increasing feelings of depression and negative self-evaluation (Stringer et al., 2010). The client may feel unsafe during a silence and avoid reporting feelings to their therapist (Levitt, 2001a). In sum, the current literature has produced varied results; silence in psychotherapy may facilitate processing of emotions or may be a strategy for clients to avoid emotional experiences. Although the psychotherapy research literature provides some information about how a client may respond emotionally to silence, there are distinct limitations to the extant literature. First, many studies arbitrarily restricted the minimum length of coded silences to between three (Frankel et al., 2006; Levitt, 2001b) and five seconds (e.g. Cuttler et al., 2019;

Implications for Practice and Policy

- Psychotherapy research is sparse on quantitative investigations of silence that include all lengths of silence during a session.
- Linguistics research theory and automated data processing techniques provide an innovative perspective to study silence in psychotherapy.
- This paper utilised theory from linguistics and automated data processing techniques to extract the full range of silences initiated by the therapist and model the client's emotion reaction in a way that has not previously been done.
- This research could help therapists understand the impact of silence and potentially contribute to training and supervision of additional therapeutic variables that impact the therapeutic process.

Sharpley et al., 2005; Daniel et al., 2018; Cook, 1964), or to 'a silent event' perceived by the therapist (Hill et al., 2003, p. 515). Therapy studies have included initial interviews (Sharpley et al., 2005), psychodynamic therapy (Hill et al., 2003), psychoanalysis (Daniel et al., 2018), general psychotherapy at a college counselling centre (Cook, 1964) and therapists providing cognitive behavioural therapy, humanistic and psychodynamic therapy (Cuttler et al., 2019). However, according to linguistics researchers, the majority of silences in everyday conversation range from 0.1 to 0.3s (Heldner & Edlund, 2010; Xiao et al., 2015; Levinson & Torreira, 2015; see Nagaoka et al., 2013, for a counterexample due to variations in language, culture and conversational norms). Silences lasting more than 3s are considered rare (see Holler et al., 2016). As such, many psychotherapy studies are likely not studying the full range of silences, and how therapeutic processes are changing from smaller silences, less than a second, to longer silences. Psychotherapy research has relied heavily on extracting silences with human coding (e.g. Pausing Inventory Categorization System—PICS—see Levitt, 2001b; Frankel et al., 2006; Daniel et al., 2018; Cuttler et al., 2019; Stringer et al., 2010; Cook, 1964), qualitative analysis (e.g. Levitt, 2001a, 2001b; Frankel & Levitt, 2009) and self-report (Hill et al., 2003; Frankel & Levitt, 2009). Although these methodologies provide some understanding of client and therapist experience of silence, they are much less temporally sensitive and likely exclude much actual inter-speaker silence that likely occurs during psychotherapy. It is possible that shorter silences also include a variety of information about the emotional nature of a session.

In addition to the restricted range of silence lengths, limited studies have examined the temporal aspect of silence and its relation to psychotherapy processes, and have produced mixed results. Cuttler et al. (2019) indicated that therapists perceived silences that facilitated emotional engagement and reflectivity as longer than silences where the client seemed disengaged, tense or confused. An analysis

of covariance in the same study also revealed that an increase in silence length corresponded to increased therapeutic collaboration. A qualitative study from Levitt (2001a) found contradictory evidence, demonstrating that clients chose to remain silent longer if they felt the therapist said something disconcerting or unpleasant. These studies show a contradiction between longer silences both indicating client disengagement and feeling unsafe, but also collaboration. These results continue to present mixed conclusions about how the length of silence is contributing to the therapeutic process. By examining only a small portion of silences, it is difficult to conclude how much silence length contributes to therapy. Alternative methodologies may be necessary to capture and study the full range of silences, and how they relate to important therapeutic processes, such as emotions.

For linguistics researchers, the study of silence fits into the larger quest for understanding how people facilitate conversation (Sacks et al., 1974; Heldner & Edlund, 2010; Levinson & Torreira, 2015). This includes how people decide who talks when (i.e. turn taking), when there are pauses in the conversation (i.e. gaps), and how individuals formulate what they are going to say (i.e. projection and reaction theory; Sacks et al., 1974; Duncan, 1972). For instance, during conversation people often anticipate when others are finished speaking and have also constructed most of what they are planning to say before their partner finishes speaking, in addition to anticipating when their partner will finish speaking (i.e. projection theory; Sacks et al., 1974; Heldner & Edlund, 2010). Speakers will use cues such as fundamental frequency (Prieto et al., 1996) to predict when their partner will finish talking. The typical distribution of silences is positively skewed, indicating that most gaps in conversation are small and that there is limited processing time post conversation exchange (for a review, see Heldner & Edlund, 2010). Using linguistics research as a model to study psychotherapy may provide additional information about how all types of silence length are related to psychological processes. For instance, Silber-Varod and Lerner (2017) compared silences ranging from 0.2 to 10s from four colloquial conversations and four therapy sessions. Results indicated that the number of between-speaker silences—one speaker stops talking, there is silence and the other speaker starts talking—was significantly different across colloquial and therapeutic conversation. Although Silber-Varod and Lerner (2017) began to explore how silence length relates to psychotherapy process, quantitative research is sparse and has yet to investigate the connections between silence length and emotional expression. Given the additional importance of emotional processing during psychotherapy (Greenberg & Safran, 1989) and the contradictory evidence regarding silence length and client emotion (Levitt, 2001a; Cuttler et al., 2019), a deeper quantitative study of all silence lengths and client emotional expression could be crucial to further understanding psychotherapy processes.

The current study presents a quantitative exploration of the silence between when the therapist stops speaking and the client starts speaking (i.e. therapist-initiated silence) during 84

psychotherapy sessions. We investigated how client emotional expression varied as therapist-initiated silences increased in length. We extracted 2744 instances of therapist-initiated silence of at least 0.02 s (Sacks et al., 1974). Emotional expression was measured via vocally encoded arousal and the emotional valence derived from semantic content (see Posner et al., 2005). Vocally encoded arousal is measured via the vocal features in the voice (see below for a further description), which are one of the most robust and minimally invasive measures of emotional activation (see Juslin & Scherer, 2005). Additionally, emotional expression from the words spoken in therapy is a thriving field of psychotherapy research and provides a minimally invasive method of extracting emotional expression (see Tanana et al., 2016; Tanana et al., 2021). We examined how both the level of client emotional expression and the variability of emotional expression related to changes in the length of silence before the client starts speaking. Given the importance that psychotherapy theorists and researchers have placed on silence, we expected that both the level and variability of client emotional expression (arousal and emotional valence) would be significantly related to silence length. However, with the variety of results in the psychotherapy literature, we were unsure as to whether emotional expression would significantly increase, potentially indicating emotional activation via emotional processing or defensiveness, or decrease, possibly indicating withdrawal or relaxing after a silence.

2 | METHODS

2.1 | Data source

For this study, we utilised a sample of 84 sessions¹ obtained from five Motivational Interviewing (MI) dissemination trials (Baer et al., 2009; Lee et al., 2014; Roy-Byrne et al., 2014; Neighbors et al., 2012; Tollison et al., 2008). At its Rogerian-based theoretical core, MI emphasises the therapist's ability to accurately understand, or make efforts to understand, a client's experience, that is, have empathy, and specifies that the therapist uses reflective statements and open questions focused on eliciting client-motivated change. Originally developed to treat individuals primarily struggling with substance use issues, MI has been expanded for use with a variety of populations (e.g. weight loss; Armstrong et al., 2011), and in many different settings (e.g. mental health, primary care; Lundahl et al., 2010). Client verbal statements about their planned, desired or actual behaviour changes (i.e. change talk) are theorised to motivate further behaviour change (see self-determination theory; Miller & Rollnick, 2012). Additionally, MI theory encourages many qualities that are considered to be therapeutically beneficial across therapies, for instance, promoting the therapeutic bond (e.g. Bordin, 1979).

Those who participated in the study were promised that no identifying information outside the recorded session would be collected. As such, there is no demographic information for the sample.

However, according to the United States Census at the time of data collection (2000), five million Washington state residents reported their race as White (84.9%), Black or African American (4%), Asian (6.7%), American Indian and Alaska Native (2.7%), Native Hawaiian and Other Pacific Islander (0.7%) or indicated another race (4.9%); in a separate measure, 7.5% indicated Hispanic or Latino ethnicity (Evans et al., 2001). Census data collected gender identification only for the gender binary (i.e. male and female) and did not include data about non-binary and transgender individuals. Given this limitation, the Census Bureau reported that 50.2% of Washington residents identified as female and 49.8% as male. Participants across studies were 18 years and older, either in a primary care setting or on a college campus.

Data from 58 therapists were utilised, all of whom conducted single-session therapy with at least one client ($M = 1.48$, $SD = 1.01$). Most therapists saw one client ($n = 44$), with three therapists seeing a maximum of five clients. The studies included therapists who completed training and received weekly supervision (Lee et al., 2013; Neighbors et al., 2012; Tollison et al., 2008), therapists who received initial training and were notified of drift from MI protocol (Roy-Byrne et al., 2014) and therapists who had received training without any continued supervision (Baer et al., 2009). In one study, primary care providers conducted either brief alcohol and drug interventions or enhanced care (Roy-Byrne et al., 2014). In three of the studies, providers targeted alcohol and marijuana use in college students (Lee et al., 2014; Neighbors et al., 2012; Tollison et al., 2008). In the last study, providers were conducting therapy in community-based primary care clinics, in which patients may have been using many types of drugs at one time (Baer et al., 2009). The primary treatment modality in each study was MI, an evidence-based treatment that emphasises empathy and specifies that the therapist uses a specific type of language (e.g. reflective statements), focused on eliciting verbal statements regarding change in a behaviour from clients (Miller & Rollnick, 2012). The trials were conducted in the Pacific-Northwest and were approved through a local institution's review board.

2.2 | Measures

2.2.1 | Silence

The definition of silence used in this research was guided by the linguistics literature, where it is established that humans can discern a pause in conversation at between 0.15 and 0.25 s (Heldner & Edlund, 2010; Levinson & Torreira, 2015), with the average being approximately 0.2 s (Campione & Véronis, 2002; Heldner & Edlund, 2010; Holler et al., 2016). Thus, we chose a silence threshold of 0.2 s; we did not restrict the maximum silence length (see Sharpley et al., 2005). Gaps of less than 0.2 s were considered 'no silence'.

The silences, as defined above, were extracted from sessions that had been manually transcribed and time stamped. This allowed for alignment between transcriptions and audio recordings, using

an Automatic Speech Recognizer (ASR) customised to the data (see Xiao et al., 2015, for further technical explanations). Sessions where alignment was not possible were discarded. An acoustic model then combed the audio data and placed timestamps when a specific phoneme was uttered. Sequences of phonemes create words, and then, words form utterances. Thus, the model composes a beginning and end of each sequence of words, and silence was defined as the space between the end of an utterance and the beginning of the next one. Transitions from one speaker to another were set at a lower threshold of 0.01 s—that is, the smallest gap in speech processed automatically where there were no overlaps (see Heldner & Edlund, 2010).

2.2.2 | Vocally encoded arousal

The first indicator of client emotional expression was arousal, measured via vocal acoustics. Vocal acoustics assist individuals to distinguish affect in social situations, which promotes effective social communication (Lima et al., 2013), and can impact the quality of social relationships, medical care and communication generally (Laukka et al., 2008). Vocal acoustics can be garnered from audio data in a variety of ways. We chose to use *Kaldi Toolkit*, a free, open-source software used for speech recognition research (available on SourceForge—<http://kaldi.sf.net/>; see Povey et al., 2011). *Kaldi* relies on two open source modelling packages, OpenFST (Allauzen et al., 2007) and two numerical algebra libraries (<http://www.netlib.org/blas/>; <http://www.netlib.org/lapack/>), in order to properly run the software. *Kaldi* first works to recognise who is speaking during a particular utterance based on unique vocal characteristics. Next, *Kaldi* transcribes a given audio file by utilising prior lexical and phonetic definitions of sounds, context of the transcript (e.g. psychotherapy data) and probability models to predict which words are likely to appear in utterances. After transcription, *Kaldi* can extract estimates of various vocal features in the audio recording (Povey et al., 2011). Using *Kaldi*, we extracted estimates of vocally encoded emotional arousal (mean f_0 ; fundamental frequency) from audio recordings of the sessions (see Juslin & Scherer, 2005; Weusthoff et al., 2013; Imel et al., 2014), every 0.025 s, with a 0.01 s gap between each estimate. Each set of mean f_0 values was assigned a speaker label (therapist or client) and put into a vector of all values for each speaker. Speaker assignment was checked by matching sections from the manually transcribed sessions to the calculated f_0 values. The median f_0 was then computed for each utterance to establish the typical arousal level for each speaker. Lastly, the median f_0 for each utterance was compared to the vector of all values, and the number of mean f_0 estimates larger than the median f_0 divided by the total number of mean f_0 estimates in a particular utterance. The resulting estimate ranged from 0 to 1. With 0.5 indicating speaker-specific average arousal, estimates greater than or equal to 0.5 indicated a higher arousal level, and estimates below 0.5 indicated a lower arousal (see Bone et al., 2014, for further technical explanation).

2.2.3 | Emotional valence

We measured positive and negative emotional expression from a metric derived from the words spoken (i.e. semantic content) by the client, which produces a continuous measure (ranging from very negative to very positive) of emotion extracted from text (Malandrakis & Narayanan, 2015). First, a dictionary of valenced words was created, and each word was weighted, known as a norm value, according to the positive and negative associations with the word (i.e. Psycholinguistic Norming). For example, the word 'happy' would have a positive valence (Turney & Littman, 2003). The basic assumption is that other words with similar meanings will have similar norm values (Malandrakis et al., 2013). The initial dictionary used was the Affective Norms for English Words (ANEW), where each word in this dictionary had a corresponding, continuous value corresponding to the magnitude (see Malandrakis & Narayanan, 2015). The weights were input into a previously researched model, which was specifically trained on psychotherapy data, and correlated with human coded words ($r = 0.87$; see Malandrakis et al., 2013). The model then computed values, ranging from -1 to 1 , on each utterance. Note that a zero-valence measure is neutral.

2.3 | Analysis

As discussed earlier, qualitative and self-report psychotherapy research has indicated limited evidence for the effects of how long a client is silent until they speak on their emotional expression, as well as evidence for variability in a client's emotional response to therapist-initiated silences. However, prior research represents only a small portion of the total silence data. We examined the changes in client emotional expression following all lengths of therapist-initiated silence with two analyses: (a) the level of emotional expression (vocally encoded arousal and emotional valence) related to silence length, and (b) variability of emotional expression in both vocally encoded arousal and emotional valence related to the length of silence.

For the analysis of level, we used a multilevel linear regression to examine the relationship between emotional expression and length of silence, with observations of silence nested within session. Silence was log transformed for the regression analysis (see Campione & Véronis, 2002). To understand variability in emotional expression, we chose to utilise an analysis of variance due to the test's methodological specificity in comparing the variance across groups. In order to create groups to compare the variance, we binned emotion into six categories to determine whether the variability in emotional expression was related to the length of silence: no silence (0–0.2s) and silences of 0.2–0.5s, 0.5–1s, 1–2s, 2–5s and greater than 5s. We first compared the six groups for mean differences in vocally encoded arousal and emotional valence with an analysis of variance (ANOVA).² We then conducted tests of variance for significant differences in vocally encoded arousal and emotional valence among the six different lengths of silence. Bartlett's test,

a type of F -test, was utilised to compare variances among multiple groups (Bartlett, 1937). Levene's test, a more robust type of F -test, was used to analyse valence due to the data skew (Schultz, 1985).

R statistical software, version 3.3.2, was used to conduct the analyses and construct the figures. Regression analysis was conducted using the *lme4* package (Bates et al., 2014). Analysis of variance and Bartlett's tests were completed with the *stats* package (Chambers et al., 2017), and Levene's test using the *car* package (Fox & Weisberg, 2019). Figures were generated with the *ggplot2* package (Wickham et al., 2019). Figures 1, 2 and 3 are density plots, generated by a kernel density estimate, which are essentially a smoothed version of a histogram.

3 | RESULTS

3.1 | Silence

As shown in Figure 1, the length of silences in this sample of psychotherapy sessions was negatively skewed, as expected (see Heldner & Edlund, 2010, Figure 3, p. 562). The mean silence was 0.96 s ($SD = 1.25$). Silences ranged between 0.2 and 24.01 s, with a median of 0.61 s and a mode of 0.23 s. On average, sessions contained 32.67 therapist-initiated silences ($SD = 21.01$) and 31.4 total s of silences ($SD = 23.6$); 98.6% of silences were under 5s, 70.7% were under 1s, and 41.4% were under 0.5s. The means, standard deviations and analytical findings for both vocally encoded arousal and emotional valence are shown in Table 1.

3.2 | Vocally Encoded Arousal

The mean client arousal across sessions was 0.56 ($SD = 0.22$). From the regression analysis, there was no significant effect of therapist-initiated silence on vocally encoded arousal ($\beta_1 = -0.01$, $t(83) = -1.71$, $p = 0.09$, 95% CI $[-0.01, 0.001]$).

Figure 2 shows the density of arousal across the six groups of silence lengths. There was a significant main effect of length of silence on mean arousal, $F(5, 4039) = 3.19$, $p < 0.01$. When examining the means in Table 1, and the plots in Figure 2, there was a decrease in mean arousal for silences greater than 5s. The effect size was small ($\eta^2 = 0.004$). Bartlett's test indicated no significant differences of arousal variance across lengths of silence ($T[5] = 1.3$, $p = 0.94$).

3.3 | Emotional Valence

The mean client valence was 0.32 ($SD = 0.13$). There was a significant effect of therapist-initiated silence on emotional valence ($\beta_1 = 0.004$, $t(83) = 2.05$, $p < 0.05$, 95% CI $[0, 0.01]$). As silence increased by 1s, emotional valence also increased by 0.004 units. Demonstrated in Figure 3, there were diminished neutral and negative responses, indicating valence became more positive as silence increased.




FIGURE 1 Density plot of silences <5s. Given that 98.6% of silences were 5 s or less, 5 s was chosen to best demonstrate the density of silence values. The number of observations greater than 5 s was 39

Figure 3 shows the density of valence across the six groups of silence lengths. There was no significant main effect of length of silence in sentiment content ($F(5, 4039) = 0.39, p = 0.86$). Levene's test demonstrated a significant difference in variance across silences of different lengths ($F(4, 4040) = 2.72, p < 0.05$). When examining Figure 3, there was a decrease in variance only for silences greater than 5 s.

4 | DISCUSSION


This study aimed to provide a large-scale temporal quantitative analysis of client emotional expression after therapist-initiated silences. Psychotherapy research has indicated a connection between silence and client emotional expression. However, prior studies have only examined longer silences, considered rare by linguistics researchers, and have produced consistently mixed results. To contribute to the existing literature, we investigated the relationship between client emotional expression and the full range of silence lengths utilising quantitative methods. Broadly, our results demonstrate a weak connection between silence length and emotional expression, indicating no persuasive evidence that silence leads to client emotional processing and expression.

To start, descriptive measures demonstrated both consistencies and discrepancies with prior linguistic and psychotherapy literatures. Consistent with linguistics findings, our results demonstrated

that the lengths of silence in psychotherapy were negatively skewed, with a modal length of silence of 0.23 s (see Heldner & Edlund, 2010). Descriptive measures indicated the vast majority (98.6%) of silences were under 5 s, consistent with prior findings that longer silences in psychotherapy are rare (Holler et al., 2016), and that many qualitative and self-report studies have explored only a small portion of silences. Cumulative statistics also indicated about 70% of silences were under 1 s, and about 40% were under 0.5 s, whereas, on average, 75% of conversational pauses are typically under 0.5 seconds (Levinson & Torreira, 2015) and between 82% and 95% are less than 1 s (Heldner & Edlund, 2010). As such, therapists may pause more often than a speaker engaged in typical colloquial conversation (see Silber-Varod & Lerner, 2017), supporting prior claims that therapists may try to be more intentional about using silence to observe the client, process therapeutic content and demonstrate their interest in the client (Hill et al., 2003).

Our hypothesis that the variance of client emotional expression would change as a function of silence length was confirmed for vocally encoded arousal; however, the effects were minimal. The ANOVA comparing arousal across groups of silence lengths indicated significant changes, but the effect size was very small. There was no significant trend in the regression analysis of arousal level changing across continuously increasing silence length. One possibility for the lack of large effects for vocally encoded arousal is our use of single-session MI data. The therapist and client are beginning to develop rapport, and not necessarily engaging in more

FIGURE 2 Emotionally encoded arousal and therapist-initiated silence



emotionally laden therapeutic material. Previous research has studied specified sessions in a series of psychotherapy (Daniel et al., 2018), as well as comparing initial and ending future sessions (Cook, 1964). Prior studies indicating emotionality and reflectivity in silence (Levitt 2001b; Hill et al., 2003) may be a product of longer and more developed therapeutic relationships (Hill et al., 2019). The current study thus provides a quantitative understanding of silence in the early phases of therapy, and in particular, the first session of therapy. Daniel et al. (2018) indicated most productive silences occurred halfway through treatment, which may indicate that client emotional expression may occur in later sessions. As such, further study may analyse longitudinal data and could explore how silence changes over time as the therapeutic relationship changes.

Our hypothesis of changing emotional valence across silence length was also confirmed, but again, the effects were weak. Regression analysis demonstrated that emotional valence significantly increased as silence length increased; however, the effect size was, again, very small. When examining Figure 3, we saw that for silences greater than 5 s, there were diminished neutral and negative responses. MI was the treatment modality, and as such, therapist goals during MI are to elicit client statements of desired change (i.e. change talk) and are discouraged from discussing statements indicating resistance to behaviour change (i.e. sustain talk) in order to promote actual behaviour change (see meta-analysis by Magill et al., 2018). It is possible that MI therapists may avoid times whereby the client is defending their behaviours, potentially eliciting

strong negative emotional reactions, leaving the existing therapeutic content to be more neutral or positive. Further research is necessary to understand how silence influences client emotional expression in other therapeutic settings.

As a quantitative and interdisciplinary study of silence, our findings are an important contribution to the psychotherapy process literature and may have some implications for clinical practice. Linguistics and computer science research provided a theoretical framework and methodology to automatically extract thousands of silences, and continuous measures of emotional expression. In addition to providing a methodology to explore the full range of silences available during a therapy session, this study also provides a method for quantitatively modelling silence and how it relates to emotional expression. Further research can expand these findings to explore how silence relates to other psychotherapy processes, as well as to address some of the study's limitations. Within a clinical context, therapists pausing more frequently than is typical in colloquial conversation supports the notion that therapy is a special type of conversation, in which traditional conversational rules and paradigms may not exist. Though clients may not necessarily be utilising silence for emotional processing, silence may create space for the client to focus on therapeutic content at their own pace (Levitt, 2002; Hill et al., 2019).

One limitation is that the study examined only therapist-initiated silences and the client's emotional response to those silences. However, pauses within a speaker's speech, as well as client-initiated




FIGURE 3 Semantic content across different lengths of silence

TABLE 1 Vocally encoded arousal and emotional valence analysis results

	Vocally encoded arousal	Emotional valence
Descriptives	Mean (SD)	Mean (SD)
Overall mean	0.56 (0.22)	0.32 (0.13)
Groups of silence		
No silence	0.58 (0.22)	0.31 (0.14)
0.2–0.5 s	0.55 (0.22)	0.31 (0.13)
0.5–1 s	0.56 (0.22)	0.32 (0.13)
1–2 s	0.56 (0.21)	0.32 (0.13)
2–5 s	0.58 (0.22)	0.32 (0.11)
Greater than 5 s	0.48 (0.21)	0.33 (0.06)
Regression analysis		
	Estimate (SE)	Estimate (SE)
Intercept	0.57 (0.01)	0.32 (0.004)
Silence coefficient (β_1)	-0.01 (0.003)	0.004 (0.002)*
95% CI	[-0.01, 0.001]	[0, 0.01]
ANOVA	$F(5, 4039) = 3.19^{**}$	$F(5, 4039) = 0.39$
Test of variance		
	Bartlett's test	Levene's test
	$T(5) = 1.3$	$F(4, 4040) = 2.72^*$

Note: Significance codes: *** $p < 0.001$; ** $p < 0.01$; * $p < 0.05$.

silences, are still relevant to explore quantitatively. Additionally, modelling the therapist's emotional expression with silence may contribute to understanding therapist-specific emotional processes, such as countertransference (see Hill et al., 1996), or client attachment style (Daniel et al., 2018). However, in terms of understanding silence as a therapist intervention in therapy, studying therapist-initiated silence may be most ideal for understanding the client's immediate emotional reaction. Future research should explore the therapist behaviours that precede silences. For instance, MI emotion process research has found that MI-inconsistent behaviours have correlated with client sustain talk (Magill et al., 2014; Pace et al., 2017; Magill et al., 2018) and that positive emotional experiences have correlated with MI-adherent behaviours (Moyers et al., 2005). However, MI process research is lacking in many ways, and, in particular, lacking research on client emotional experiences (Wagner & Ingersoll, 2008).

The current data also lack demographic information to better understand the relationship between individuals' identities and silence (e.g. gender; Ramakrishna et al., 2015), and a broader relationship of silence, cultural norms (Nagaoka et al., 2013) and relational power dynamics. Studies have primarily examined silences in the context of European countries and the mainland and continental United States (Heldner & Edlund, 2010). Future research could benefit from including demographic variables, therapeutic process

variables and therapy outcomes. Even with these limitations, the study provides a strong basis for quantitative and interdisciplinary research on silence during psychotherapy and contributes to the understanding of how smaller exchanges between therapist and client contribute to larger psychotherapy processes.

ORCID

Christina S. Soma  <https://orcid.org/0000-0002-3607-6643>

ENDNOTES

- ¹ The full data set included both real and standardised patients. We included only real patients in this sample. Additionally, the 84 sessions were able to be processed for silence with our data processing procedure. Due to the nature of community mental health, the quality of some audio recordings was not conducive to extracting silence with our methodology.
- ² Due to the nested nature of psychotherapy data, ANOVA and tests of variance compared the entire data set, as well as a random sample of data containing one client per therapist to control for nesting. Results were consistent in both sets of tests.

REFERENCES

- Ackerman, K. D., Martino, M., Heyman, R., Moyna, N. M., & Rabin, B. S. (1998). Stressor-induced alteration of cytokine production in multiple sclerosis patients and controls. *Psychosomatic Medicine*, *60*(4), 484–491.
- Allauzen, C., Riley, M., Schalkwyk, J., Skut, W., & Mohri, M. (2007). OpenFst: A general and efficient weighted finite-state transducer library. In *International Conference on Implementation and Application of Automata* (pp. 11–23). Springer.
- Armstrong, M. J., Mottershead, T. A., Ronksley, P. E., Sigal, R. J., Campbell, T. S., & Hemmelgarn, B. R. (2011). Motivational interviewing to improve weight loss in overweight and/or obese patients: A systematic review and meta-analysis of randomized controlled trials. *Obesity Reviews*, *12*(9), 709–723.
- Baer, J. S., Wells, E. A., Rosengren, D. B., Hartzler, B., Beadnell, B., & Dunn, C. (2009). Agency context and tailored training in technology transfer: A pilot evaluation of motivational interviewing training for community counselors. *Journal of Substance Abuse Treatment*, *37*, 191–202.
- Bar-On, R., Tranel, D., Denburg, N. L., & Bechara, A. (2004). Emotional and social intelligence. *Social Neuroscience: Key Readings*, 223.
- Bartlett, M. S. (1937). Properties of sufficiency and statistical tests. *Proceedings of the Royal Society of London. Series A: Mathematical and Physical Sciences*, *160*(901), 268–282.
- Bates, D., Mächler, M., Bolker, B., & Walker, S. (2014). Fitting linear mixed-effects models using lme4. *arXiv preprint arXiv:1406.5823*.
- Bone, D., Lee, C. C., & Narayanan, S. (2014). Robust unsupervised arousal rating: A rule-based framework with knowledge-inspired vocal features. *IEEE Transactions on Affective Computing*, *5*(2), 201–213.
- Bordin, E. S. (1979). The generalizability of the psychoanalytic concept of the working alliance. *Psychotherapy: Theory, research & practice*, *16*(3), 252.
- Campione, E., & Véronis, J. (2002). A large-scale multilingual study of silent pause duration. In *Speech prosody 2002, international conference*.
- Chambers, J. M., Freeny, A. E., & Heiberger, R. M. (2017). Analysis of variance; designed experiments. In *Statistical models in S* (pp. 145–193). Routledge.
- Cook, J. J. (1964). Silence in psychotherapy. *Journal of Counseling Psychology*, *11*(1), 42.
- Cuttler, E., Hill, C. E., King, S., & Kivlighan, D. M., Jr. (2019). Productive silence is golden: Predicting changes in client collaboration from process during silence and client attachment style in psychodynamic psychotherapy. *Psychotherapy*, *56*(4), 568.
- Daniel, S. I., Folke, S., Lunn, S., Gondan, M., & Poulsen, S. (2018). Mind the gap: In-session silences are associated with client attachment insecurity, therapeutic alliance, and treatment outcome. *Psychotherapy Research*, *28*(2), 203–216.
- Duncan, S. (1972). Some signals and rules for taking speaking turns in conversations. *Journal of Personality and Social Psychology*, *23*(2), 283.
- Evans, D., Price, J., & Barron, W. (2001). *Profiles of general demographic characteristics: 2000 Census of population and housing*. Washington, DC: US Department of Commerce.
- Fox, J., & Weisberg, S. (2019). *An R Companion to Applied Regression* (Third ed.). Sage.
- Frank, J. D. (1961). *Persuasion and Healing*. The Johns Hopkins University Press.
- Frankel, Z. E., & Levitt, H. M. (2009). Clients' experiences of disengaged moments in psychotherapy: A grounded theory analysis. *Journal of Contemporary Psychotherapy*, *39*(3), 171–186.
- Frankel, Z. E., Levitt, H. M., Murray, D. M., Greenberg, L. S., & Angus, L. (2006). Assessing silent processes in psychotherapy: An empirically derived categorization system and sampling strategy. *Psychotherapy Research*, *16*(5), 627–638.
- Freud, S. (1912). *Recommendations to physicians practising psycho-analysis*. The.
- Gale, J., & Sanchez, B. (2005). The meaning and function of silence in psychotherapy with particular reference to a therapeutic community treatment programme. *Psychoanalytic Psychotherapy*, *19*(3), 205–220.
- Gladstein, G. A. (1974). Nonverbal communication and counseling/psychotherapy: A review. *The Counseling Psychologist*, *4*(3), 34–57.
- Greenberg, L. S., & Safran, J. D. (1989). Emotion in psychotherapy. *American Psychologist*, *44*(1), 19.
- Heldner, M., & Edlund, J. (2010). Pauses, gaps and overlaps in conversations. *Journal of Phonetics*, *38*(4), 555–568.
- Helm, J. L., Sbarra, D. A., & Ferrer, E. (2014). Coregulation of respiratory sinus arrhythmia in adult romantic partners. *Emotion*, *14*(3), 522.
- Hill, C. E., Nutt-Williams, E., Heaton, K. J., Thompson, B. J., & Rhodes, R. H. (1996). Therapist retrospective recall impasses in long-term psychotherapy: A qualitative analysis. *Journal of Counseling Psychology*, *43*(2), 207.
- Hill, C. E., Thompson, B. J., & Ladany, N. (2003). Therapist use of silence in therapy: A survey. *Journal of Clinical Psychology*, *59*(4), 513–524.
- Hill, C. E., Kline, K. V., O'Connor, S., Morales, K., Li, X., Kivlighan, D. M., Jr., & Hillman, J. (2019). Silence is golden: A mixed methods investigation of silence in one case of psychodynamic psychotherapy. *Psychotherapy*, *56*(4), 577.
- Holler, J., Casillas, M., Kendrick, H., & K., & C Levinson, S. (2016). *Turn-taking in human communicative interaction*. Frontiers Media SA.
- Imel, Z. E., Barco, J. S., Brown, H. J., Baucom, B. R., Kircher, J. C., Baer, J. S., & Atkins, D. C. (2014). The Association of Therapist Empathy and Synchrony in Vocally Encoded Arousal. *Journal of Counseling Psychology*, *61*(1), 146–153.
- Isen, A. M. (2008). Some ways in which positive affect influences decision making and problem solving. *Handbook of Emotions*, *3*, 548–573.
- Johannesen, R. L. (1974). The functions of silence: A plea for communication research. *Western Journal of Communication (includes Communication Reports)*, *38*(1), 25–35.
- Juslin, P. N., & Scherer, K. R. (2005). *Vocal expression of affect*. Oxford University Press.
- Kiecolt-Glaser, J. K., Loving, T. J., Stowell, J. R., Malarkey, W. B., Lemeshow, S., Dickinson, S. L., & Glaser, R. (2005). Hostile marital interactions, proinflammatory cytokine production, and wound healing. *Archives of General Psychiatry*, *62*(12), 1377–1384.

- Ladany, N., Hill, C. E., Thompson, B. J., & O'Brien, K. M. (2004). Therapist perspectives on using silence in therapy: A qualitative study. *Counselling and Psychotherapy Research, 4*(1), 80–89.
- Lambert, M. J., & Bergin, G. (2004). Overview, Trends and Future Issues. In M. J. Lambert (Ed.), *Bergin and Garfield's Handbook of Psychotherapy and Behaviour Change*. John Wiley & Sons.
- Lane, R. C., Koetting, M. G., & Bishop, J. (2002). Silence as communication in psychodynamic psychotherapy. *Clinical Psychology Review, 22*(7), 1091–1104.
- Lang, P. J., & Bradley, M. M. (2010). Emotion and the motivational brain. *Biological Psychology, 84*(3), 437–450.
- Laukka, P., Linnman, C., Åhs, F., Pissioti, A., Frans, Ö., Faria, V., ... Furmark, T. (2008). In a nervous voice: Acoustic analysis and perception of anxiety in social phobics' speech. *Journal of Nonverbal Behaviour, 32*(4), 195.
- Lee, C. M., Neighbors, C., Lewis, M. A., Kaysen, D., Mittman, A., Geisner, I. M., Atkins, D. C., Zheng, C., Garberson, L. A., Kilmer, J. R., & Larimer, M. E. (2014). Randomized Controlled Trial of a Spring Break Intervention to Reduce High-Risk Drinking. *Journal of Counseling and Clinical Psychology, 82*(2), 189–201.
- Lee, C. M., Kilmer, J. R., Neighbors, C., Atkins, D. C., Zheng, C., Walker, D. D., & Larimer, M. E. (2013). Indicated prevention for college student marijuana use: A randomized controlled trial. *Journal of Counseling and Clinical Psychology, 81*(4), 702–709.
- Levinson, S. C., & Torreira, F. (2015). Timing in turn-taking and its implications for processing models of language. *Frontiers in Psychology, 6*, 731.
- Levitt, H. M. (2001a). Clients' experiences of obstructive silence: Integrating conscious reports and analytic theories. *Journal of Contemporary Psychotherapy, 31*(4), 221–244.
- Levitt, H. M. (2001b). Sounds of silence in psychotherapy: The categorization of clients' pauses. *Psychotherapy Research, 11*(3), 295–309.
- Levitt, H. M. (2002). The unsaid in the psychotherapy narrative: Voicing the unvoiced. *Counselling Psychology Quarterly, 15*(4), 333–350.
- Lima, C. F., Castro, S. L., & Scott, S. K. (2013). When voices get emotional: A corpus of nonverbal vocalizations for research on emotion processing. *Behaviour Research Methods, 45*(4), 1234–1245.
- Lundahl, B. W., Kunz, C., Brownell, C., Tollefson, D., & Burke, B. L. (2010). A meta-analysis of motivational interviewing: Twenty-five years of empirical studies. *Research on Social Work Practice, 20*(2), 137–160.
- Magill, M., Gaume, J., Apodaca, T. R., Walthers, J., Mastroleo, N. R., Borsari, B., & Longabaugh, R. (2014). The technical hypothesis of motivational interviewing: a meta-analysis of MI's key causal model. *Journal of Consulting and Clinical Psychology, 82*(6), 973–983.
- Magill, N., Graves, H., de Zoysa, N., Winkley, K., Amiel, S., Shuttlewood, E., Landau, S., & Ismail, K. (2018). Assessing treatment fidelity and contamination in a cluster randomised controlled trial of motivational interviewing and cognitive behavioural therapy skills in type 2 diabetes. *BMC Family Practice, 19*(1), 60.
- Malandrakis, N., & Narayanan, S. S. (2015). Therapy language analysis using automatically generated psycholinguistic norms. In the Sixteenth Annual Conference of the International Speech Communication Association.
- Malandrakis, N., Potamianos, A., Iosif, E., & Narayanan, S. (2013). Distributional semantic models for affective text analysis. *IEEE Transactions on Audio, Speech and Language Processing, 21*(11), 2379–2392.
- Matarazzo, J. D., & Wiens, A. N. (1977). Speech behaviour as an objective correlate of empathy and outcome in interview and psychotherapy research: A review with implications for behaviour modification. *Behaviour Modification, 1*(4), 453–480.
- Miller, W. R., & Rollnick, S. (2012). *Motivational interviewing: Helping people change*. Guilford press.
- Moyers, T. B., Miller, W. R., & Hendrickson, S. M. L. (2005). How does motivational interviewing work? Therapist interpersonal skill predicts client involvement within motivational interviewing sessions. *Journal of Consulting and Clinical Psychology, 73*(4), 590–598.
- Nagaoka, C., Kuwabara, T., Yoshikawa, S., Watabe, M., Komori, M., Oyama, Y., & Hatanaka, C. (2013). Implication of silence in a Japanese psychotherapy context: A preliminary study using quantitative analysis of silence and utterance of a therapist and a client. *Asia Pacific Journal of Counselling and Psychotherapy, 4*(2), 147–152.
- Neighbors, C., Lee, C. M., Atkins, D. C., Lewis, M. A., Kaysen, D., Mittman, A., Fossos, N., Geisner, I. M., Zheng, C., Garberson, L. A., Kilmer, J. R., & Larimer, M. E. (2012). A Randomized Controlled Trial of Event-Specific Prevention Strategies for Reducing Problematic Drinking Associated with 21st Birthday Celebrations. *Journal of Counseling and Clinical Psychology, 90*(5), 850–862.
- Pace, B. T., Dembe, A., Soma, C. S., Baldwin, S. A., Atkins, D. C., & Imel, Z. E. (2017). A multivariate meta-analysis of motivational interviewing process and outcome. *Psychology of Addictive Behaviours: Journal of the Society of Psychologists in Addictive Behaviours, 31*(5), 524–533.
- Picard, M. (1952). *The World of Silence*. Regnery.
- Posner, J., Russell, J. A., & Peterson, B. S. (2005). The circumplex model of affect: An integrative approach to affective neuroscience, cognitive development, and psychopathology. *Development and Psychopathology, 17*(3), 715.
- Povey, D., Ghoshal, A., Boulianne, G., Burget, L., Glembek, O., Goel, N., ... Silovsky, J. (2011). The Kaldi speech recognition toolkit. In *IEEE 2011 workshop on automatic speech recognition and understanding* (No. CONF). IEEE Signal Processing Society.
- Prieto, P., Shih, C., & Nibert, H. (1996). Pitch downtrend in Spanish. *Journal of Phonetics, 24*, 445–473.
- Ramakrishna, A., Malandrakis, N., Staruk, E., & Narayanan, S. (2015). A quantitative analysis of gender differences in movies using psycholinguistic normatives. In *Proceedings of the 2015 Conference on Empirical Methods in Natural Language Processing* (pp. 1996–2001).
- Roy-Byrne, P., Bumgardner, K., Krupski, A., Dunn, C., Ries, R., Donovan, D., ... Joesch, J. M. (2014). Brief intervention for problem drug use in safety-net primary care settings: a randomized clinical trial. *JAMA, 312*(5), 492–501.
- Sacks, H., Schegloff, E. A., & Jefferson, G. (1974). A simplest systematics for the organization of turn-taking for conversation. *Language, 50*, 696–735.
- Safran, J. D., & Muran, J. C. (2000). Resolving therapeutic alliance ruptures: Diversity and integration. *Journal of Clinical Psychology, 56*(2), 233–243.
- Sharpley, C. F., Munro, D. M., & Elly, M. J. (2005). Silence and rapport during initial interviews. *Counselling Psychology Quarterly, 18*(2), 149–159.
- Schultz, B. B. (1985). Levene's test for relative variation. *Systematic Zoology, 34*(4), 449–456.
- Silber-Varod, V., & Lerner, A. (2017). Analysis of silences in unbalanced dialogues: the effect of genre and role. *TMH-QPSR, 53*.
- Stringer, J. V., Levitt, H. M., Berman, J. S., & Mathews, S. S. (2010). A study of silent disengagement and distressing emotion in psychotherapy. *Psychotherapy Research, 20*(5), 495–510.
- Tanana, M., Dembe, A., Soma, C. S., Imel, Z., Atkins, D., & Srikumar, V. (2016, June). Is sentiment in movies the same as sentiment in psychotherapy? comparisons using a new psychotherapy sentiment database. In *Proceedings of the Third Workshop on Computational Linguistics and Clinical Psychology* (pp. 33–41).
- Tanana, M. J., Soma, C. S., Kuo, P. B., Bertagnolli, N. M., Dembe, A., Pace, B. T., ... Imel, Z. E. (2021). How do you feel? Using natural language processing to automatically rate emotion in psychotherapy. *Behaviour Research Methods, 53*(5), 2069–2082.
- Tollison, S. J., Lee, C. M., Neighbors, C., Neil, T. A., Olson, N. D., & Larimer, M. E. (2008). Questions and Reflections: The Use of Motivational Interviewing Microskills in a Peer-Led Brief Alcohol Intervention for College Students. *Behaviour Therapy, 39*(2), 183–194.

- Turney, P. D., & Littman, M. L. (2003). Measuring praise and criticism: Inference of semantic orientation from association. *ACM Transactions on Information Systems (TOIS)*, 21(4), 315–346.
- Wagner, C. C., & Ingersoll, K. S. (2008). Beyond Cognition: Broadening the Emotional Base of Motivational Interviewing. *Journal of Psychotherapy Integration*, 18(2), 191–206.
- Weusthoff, S., Baucom, B. R., & Hahlweg, K., (2013). The siren song of vocal fundamental frequency for romantic relationships. *Frontiers in Psychology*, 4, 439.
- Wickham, H., Averick, M., Bryan, J., Chang, W., McGowan, L., François, R., ... Kuhn, M. (2019). Welcome to the Tidyverse. *Journal of Open Source Software*, 4(43), 1686.
- Xiao, B., Imel, Z. E., Atkins, D. C., Georgiou, P. G., & Narayanan, S. S. (2015). Analyzing speech rate entrainment and its relation to therapist empathy in drug addiction counseling. In *INTERSPEECH* (pp. 2489–2493).

AUTHOR BIOGRAPHIES

Christina S. Soma is a PhD candidate at the University of Utah, working with Dr. Zac Imel. She recently returned from a research fellowship at the Modum Bad Psychiatric Hospital in Vikersund, Norway. She researches how the little moments in psychotherapy contribute to the broader therapist-client process. She's currently a doctoral psychology intern at the Colorado State University Health Network, and will soon start as a Postdoctoral Fellow at Lyssn, Inc.

Bruce E. Wampold, Ph.D., is professor emeritus at the University of Wisconsin-Madison and Chief Scientist, SkillsSetter.com. He has studied the effectiveness of psychotherapy, therapist effects, and the characteristics and actions of effective therapists. He has also proposed a Contextual Model to explain the benefits of psychotherapy.

Nikolaos Flemotomos is a PhD candidate in Electrical Engineering, and member of the Signal Analysis and Interpretation Laboratory at the University of Southern California (USC). His research interests include machine learning, speech, and language processing focused on structured conversational interactions. He holds an undergraduate degree in Electrical and Computer Engineering from the National Technical University of Athens (Greece) and MSc degrees from USC in Electrical Engineering and in Computer Science. He has worked as a research intern with the Czech Technical University in Prague, with Microsoft Research and with Apple on multiple sub-fields of signal and speech processing.

Raghuveer Peri (Member, IEEE) is a PhD candidate in the Electrical and Computer Engineering Department, University of Southern California, Los Angeles, CA, USA. Prior to this, he was a Research Engineer with Multimedia advanced R&D Team, Qualcomm, San Diego, CA, USA. His current research interests include robust speaker verification, speaker diarization, and adversarial attacks on speaker identification systems.

Shrikant Narayanan is University Professor and Niki & C. L. Max Nikias Chair in Engineering at the University of Southern California, and holds appointments as Professor of Electrical and Computer Engineering, Computer Science, Linguistics, Psychology, Neuroscience, Otolaryngology-Head and Neck Surgery, and Pediatrics, Research Director of the Information Science Institute, and director of the Ming Hsieh Institute. His research focuses on developing engineering approaches to understand the human condition and in creating machine intelligence technologies that can support and enhance human experiences. He is a Fellow of the Acoustical Society of America, IEEE, ISCA, the American Association for the Advancement of Science, the Association for Psychological Science, the American Institute for Medical and Biological Engineering and the National Academy of Inventors.

David C. Atkins, Ph.D., is Research Professor of Psychiatry and Behavioral Sciences at the University of Washington and core faculty of the Behavioral Research in Technology and Engineering (BRiTE) Center, focused on technology and mental health. He leads an interdisciplinary research team including engineers, computer scientists, designers, and clinical researchers who develop spoken language technologies to estimate quality metrics in counseling, and how such technologies can assist training, supervision, and quality assurance of evidence-based counseling services. In addition to his academic work, Dr. Atkins is a co-founder of a start-up, Lyssn.io, that is focused on developing and implementing technology to support evidence-based counseling.

Zac E. Imel is a Professor with the Counseling Psychology Program in the Department of Educational Psychology and Adjunct Assistant Professor in the Department of Psychiatry at the University of Utah. His primary interests involve research, teaching, and service related to the promotion and understanding of quality mental health treatment. Specific programs of research include methods for identifying and understanding the behaviors of effective (and less effective) therapists, the utilization of mental health services, emerging linguistic techniques for modeling psychotherapy process, and meta-analysis of treatment outcome studies.

How to cite this article: Soma, C. S., Wampold, B. E., Flemotomos, N., Peri, R., Narayanan, S., Atkins, D. C., & Imel, Z. E. (2022). The Silent Treatment? Changes in patient emotional expression after silence. *Counselling and Psychotherapy Research*, 00, 1–11. <https://doi.org/10.1002/capr.12537>